

Advertise

Subscribe



HOME ABOUT CONTINUING EDUCATION ARTICLES INDUSTRY NEWS WEBINARS

LOG IN FOR CE CREDITS

# The use of dynamic navigation to prevent implant complications — every patient, every time

APRIL 22, 2017 BY Implant Practice Team

1 0 Tweet 0

## Drs. Robert W. Emery and Keith Progebin discuss navigation for accurate and precise implant planning

Complications related to dental implants fall into two broad categories: infectious and mechanical. The underlying etiology of these two categories often results from poor implant position.<sup>1-3</sup> Computer-assisted surgery (CAS) allows the dental team, including restorative dentist, surgeon, and dental technician, to plan and place implants more accurately and precisely. There are two forms of CAS:

- Static guidance, which uses prefabricated CAD/CAM splints, proprietary tubes, and drills to guide the surgeon’s instruments. These guides cannot be altered after fabrication and are thus “static.”
- Dynamic guidance, which uses a stereotactic tracking system to dynamically guide the surgeon’s instruments to the correct location. The plan can be altered at any time and is thus “dynamic.”


With the advent of dynamic navigation for the placement of dental implants, the dental surgeon can place implants accurately and precisely every time in an efficient and cost-effective fashion.<sup>4</sup> A complete digital workflow used with dynamic guidance requires no lab work. The patient can be scanned, the case planned, and the implant placed in one visit. The typical cost for a dynamically

f G+ in



Search this website

Promotions



Promotions

guided case is less than \$100. Presently, dental surgeons often guide only the “tough” or “complex” cases. This “complex-only” approach has resulted from the difficulty and cost of using static guides. Static guides require that the guides be fabricated ahead of time. Once fabricated, the guides cannot be changed. The cost varies from \$300 to \$1,200 per case.

The literature clearly shows that dentists are not as accurate and precise when they place implants freehand compared with any form of guided surgery.<sup>5–8</sup> This case study illustrates the management of a significant complication related to the placement of a single implant in the wrong position. The “simple” becomes “complex.”

This 45-year-old male presented to the surgeon with a complaint of a failed implant and loose adjacent teeth. The implant had been placed by another surgeon 3 months earlier. Initially, the patient was told he had a vertically fractured tooth No. 4 that was not restorable (Figure 1). The tooth was extracted and an immediate implant placed freehand (Figure 2). The patient described pain and tenderness that persisted following implant placement. The patient went back to the surgeon on a weekly basis and received two-dimensional radiographs. The postoperative course included an infection treated with antibiotics for 1 month and eventual removal of the implant No. 4. The patient now had tenderness around the implant site with mobility and deep periodontal pockets on teeth Nos. 3 and 5. Cone beam computed tomography (CBCT) revealed periapical abscesses of teeth Nos. 3 and 5 with loss of adjacent periodontal attachments (Figure 3).



Figure 1: Pre-surgical periapical, tooth No. 4 fractured vertically




Figure 2: Postoperative periapical showing incorrect implant position with impingement of periodontal ligament of tooth No. 3

**Unlock**  
your access to  
**24 CE credits**

Digital edition subscribers have  
access to the full CE library...

Subscribe Today!



**Implant**  
Practice US

## Sign Up For E-News

Join our E-News for updates on great articles from Implant Practice US, as well as upcoming events and opportunities for CE credits.

First Name

Last Name

Email Address

**SIGN UP!**

Read the Latest  
Issue!




Figure 3: CBCT showing periapical abscess and attachment loss of teeth Nos. 3 and 5

The patient was referred for prosthetic consultation, and a treatment plan was developed. The plan included the following:

1. Intraoral scanning for virtual treatment planning. Diagnostic wax-up and fabrication of a removable provisional prosthesis for areas Nos. 3, 4, 5 (Essix retainer)
2. Extraction of teeth Nos. 3 and 5 with immediate guided-tissue regeneration using cortico-cancellous allograft, PRF, and dense PTFE membrane
3. Allow the area to heal for 2 months
4. Placement of implant Nos. 3, 4, and 5 using dynamic navigation (X-Guide, X-Nav Technologies, LLC, Lansdale, Pennsylvania) and osseodensification (Densah<sup>®</sup>, Versah, LLC, Jackson, Michigan)
5. Allow 4 months for osseointegration followed by ISQ evaluation
6. Provisionalization for 2 to 4 months
7. Final fixed restoration

The patient underwent extraction and immediate guided-tissue regeneration (Figures 4 and 5). A removable provisional restoration interim appliance was placed with soft tissue conditioner, and the graft then matured for 2 months.



Like Us On  
Facebook



Latest Tweets



Figure 4: Surgical view at the time of extraction and immediate guided-tissue reconstruction. Note the loss of adjacent periodontal attachment




Figure 5: The immediate post-surgical panoramic




Figure 6: Post-op CBCT (dark blue) superimposed on pre-surgical plan (light blue) to illustrate accuracy of dynamic image navigation

At 2 months, the patient underwent pre-surgical CBCT with an X-Clip. The DICOM of the CBCT was imported in the X-Guide planning software and the pre-extraction stl file imported for planning purposes. The implants were dynamically guided into position using the X-Guide. The post-surgical CBCT was superimposed on the pre-surgical plan to evaluate the accuracy of implant placement (Figure 6). The accuracy (mean) angular deviation from the plan for all three implants was 4.2 degrees. The precision (standard deviation) was 1.6 degrees. The accuracy (mean) platform entry deviation was 0.5 mm, and the precision (standard deviation) was 0.36 mm. After a period of 4 months, ISQ values greater than 70 were found for all three implants. A fixed provisional restoration was fabricated to manipulate the tissue, transitionally load the implants, and develop an appropriate hygiene regime for the patient (Figure 7). Only after the patient's confidence and expectations were met was the definitive prosthesis fabricated and then restored.

Tweets by  
@ImplantPractice

 **Implant Practic...**  
@ImplantPractice

Carbon coating can reduce heat and vibrations while using Burs. #excalibur

[implant-us.link/2rJu61o](https://implant-us.link/2rJu61o)



34m

 **Implant Practic...**  
@ImplantPractice

Trainer Joel Freeman leading workout at Dentsply Sirona World 2017 [implant-us.link/2sTcpMN](https://implant-us.link/2sTcpMN)



[Embed](#) [View on Twitter](#)

The patient is currently on a 3-month maintenance program to reevaluate his oral hygiene, restore his confidence in dentistry, and ensure a successful outcome. The final splinted ceramometal bridge was cemented with Temrex cement on gold custom abutments. (Figure 8).




Figure 7: Fixed splinted provisional acrylic-cemented restorations



Figure 8: Fixed ceramometal bridge cemented on gold custom abutments Nos. 3I-4I-5I

Prevention is the best way to avoid complications. Published data has shown that implants placed freehand are statistically less precise and accurate in every measure.<sup>5-7,9,10</sup> Block showed freehand mean angular deviation was 6.5 degrees with a standard deviation of 4.21 degrees, compared to dynamically guided implants with a mean angular deviation of 2.9 degrees with a standard deviation of 1.36 degrees for fully guided.<sup>5</sup> This case illustrates the time efficient digital workflow with no pre-surgical laboratory work necessary to implement dynamic

guidance. The efficient workflow and decreased costs relative to static guidance allow this form of guidance to be used on every patient, every time. Had this patient had this “simple” case performed with any form of guidance, this “complex” complication could have been avoided.

In summary, dynamic navigation has given the dental team the ability to immediately and efficiently scan, plan, coordinate our care, and place implants more accurately and precisely on every patient, every time, thus avoiding complications.

## Author Info



Robert W. Emery, BDS, DDS, is a board-certified Oral and Maxillofacial Surgeon in private practice in Washington, D.C. Dr. Emery received the Nobelpharma Oral and Maxillofacial Surgery Research Award and Grant from the Oral and Maxillofacial Surgery Research Foundation. He is actively involved in numerous research projects with an emphasis on image-guided surgery and is a founding partner of X-Nav Technologies, LLC. Presently, he is Director of the Capital Center for Oral and Maxillofacial Surgery, Senior Attending Surgeon at the Washington Hospital Center, and Chief Medical Officer of X-Nav Technologies.



Keith Progebin, DDS, is a former Associate Clinical Instructor, Department of Prosthodontics and Occlusion, New York University College of Dentistry. Dr. Progebin is a Member of the Academy of Osseointegration and the American College of Prosthodontics. He is also a Fellow of the Greater New York Academy of Prosthodontics, the International College of Dentists, and the American College of Dentists. Dr. Progebin is in Private Practice in Advanced Restorative Dentistry and Implant Prosthodontics in Washington, D.C.

## References

1. Link-Bindo EE, Soltys J, Donatelli D, Cavanaugh R. Common Prosthetic Implant Complications in Fixed Restorations. *Compend Contin Educ Dent*. 2016;37(7):431-436.

2. Roos-Jansåker AM. Long time follow up of implant therapy and treatment of peri-implantitis. *Swed Dent J Suppl.* 2007;188:7-66.
3. Shim HW, Yang BE. Long-term cumulative survival and mechanical complications of single-tooth Ankylos Implants: focus on the abutment neck fractures. *J Adv Prosthodont.* 2015;7(6):423-430.
4. Block MS, Emery RW. Static or Dynamic Navigation for Implant Placement — Choosing the Method of Guidance. *J Oral Maxillofac Surg.* 2016;74(2):269-277.
5. Block MS, Emery RW, Cullum DR, Sheikh A. Implant Placement is more accurate using dynamic navigation. *J Oral Maxillofac Surg.* 2017. Article in Press. [http://www.joms.org/article/S0278-2391\(17\)30252-5/fulltext?rss=yes](http://www.joms.org/article/S0278-2391(17)30252-5/fulltext?rss=yes). Accessed March 21, 2017.
6. Block MS, Emery RW, Lank K, Ryan J. Implant Placement Accuracy Using Dynamic Navigation. *Int J Oral Maxillofac Implants.* 2017;32(1):92-99.
7. Emery RW, Merritt SA, Lank K, Gibbs JD: Accuracy of Dynamic Navigation for Dental Implant Placement—Model-Based Evaluation. *J Oral Implantol.* 2016;42(5):399-405.
8. Tahmaseb A, Wismeijer D, Coucke W, Derksen DW. Computer Technology Applications in Surgical Implant Dentistry: A systematic Review. *Int J Oral Maxillofac Implants.* 2014;29(Suppl):25-42.
9. Vercruyssen M, Cox C, Coucke W, Naert I, Jacobs R, Quirynen M: A randomized clinical trial comparing guided implant surgery (bone- or mucosa-supported) with mental navigation or the use of a pilot-drill template. *J Clin Periodontol.* 2014;41(7):717-723.
10. Kramer FJ, Baethge C, Swennen G, Rosahl S. Navigated vs. conventional implant insertion for maxillary single tooth replacement. *Clin Oral Implants Res.* 2005;16(1):60-68.

FILED UNDER: [Current Issue](#), [Technology](#)

---

## Our Magazine

[Read Current Issue](#)

[About](#)

[Our Team](#)

[Advertise With Us](#)

[About Us](#)

[Contact Us](#)

[Privacy Policy](#)

[Terms & Conditions](#)


## Subscribe Today

Subscribe to Implant Practice US today online: get access to 24 CE Credits per year. Click the link below to learn more about our subscription options.

[Subscribe](#)

## CE Credits

Log in for CE Credits



MedMark, LLC is the leading interactive marketing and advertising company specializing in medical advertising, custom media and public relations for the U.S. dental industry.